



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\angle OAG = \angle ODH. \therefore GA = DH. \text{ Similarly } CG = BH.$$

$$\therefore AG + BH = CG + DH.$$

$$\therefore \frac{1}{2}h(AG + BH) = \frac{1}{2}h(CG + DH).$$

$$\therefore \text{Area } AGHD = \text{area } CGHD.$$

Good solutions of this problem were received from *Professors Wm. Symmonds, and Cooper D. Schmitt.*

This problem has proved to be a very interesting one and for that reason we have given it extra space. The proposition to which Prof. Ellwood has given a proof explanatory to the proposition under consideration is known as the proposition of *Menelaus*. See *Halsted's Elementary Synthetic Geometry*, p. 117. Editor.

PROBLEMS.

42. Proposed by G. I. HOPKINS, Instructor in Mathematics and Physics in High School, Manchester, New Hampshire.

If the bisectors of two angles of a triangle are equal the triangle is isosceles.

[The term *bisector* in this theorem means the line which divides an angle into two equal parts and terminates in the opposite side.]

43. Proposed by J. F. W. SCHEFFER, Hagerstown, Maryland.

The consecutive sides of a quadrilateral are a, b, c, d . Supposing its diagonals to be equal, find them and also the area of the quadrilateral.

CALCULUS.

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

27. Proposed by Professor G. B. M. ZERR, A. M., Principal of High School, Staunton, Virginia.

A runs around the circumference of a circular field with velocity m feet; B starts from the centre with velocity $n > m$ feet to catch A . The straight line joining their positions always passes through the centre. Find the equation to the curve described by B , the distance he runs and the time occupied.

L. Solution by A. H. HOLMES, Brunswick, Maine, H. W. DRAUGHON, Ohio, Mississippi, and the PROPOSER.

Let A be the point of starting of the pursued, P, B , the position of the pursuer and pursued at any time.

$$\text{Let } OA = a, OP = r, \angle BOA = \theta, \frac{m}{n} = u, \text{ arc } OP = s.$$